

REPORT D

AD-A240 378

Form Approved  
OMB No. 0704-0188

Public reporting burden for this collection of information is  
maintaining the data needed, and completing and reviewing  
suggestions for reducing this burden, to Washington Headq  
and to the Office of Management and Budget, Paperwork



Instructions, searching existing data sources, gathering and  
or any other aspect of this collection of information, including  
erson Davis Highway, Suite 1204, Arlington, VA 22202-4302.

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE August 1991		3. REPORT TYPE AND DATES COVERED Professional paper	
4. TITLE AND SUBTITLE CONDITIONAL EVENT ALGEBRAS: TWO NEW CHARACTERIZATIONS AND THEIR RELATIONS TO BAYESIAN ANALYSIS				5. FUNDING NUMBERS PR: ZT52 PE: 0601152N WU: DN306225	
6. AUTHOR(S) I. R. Goodman				8. PERFORMING ORGANIZATION REPORT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Naval Ocean Systems Center San Diego, CA 92152-5000					
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) Office of Chief of Naval Research Independent Research Programs (IR) OCNR-10P Arlington, VA 22217-5000				10. SPONSORING/MONITORING AGENCY REPORT NUMBER	
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION/AVAILABILITY STATEMENT  Approved for public release; distribution is unlimited.				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words)  The standard approach in developing conditional probability is a numerically oriented one, although one notable exception has been efforts in the area of qualitative (comparative-preference ordering) probability theory, as developed by Suppes, Domotor, Fishburn, and others. However, the latter aspect of probability requires, in general, the preference ordering to be equivalent to the probability ordering. But, this is obviously too restrictive to be compatible with the basic monotonicity property of probability, relative to the usual partial subclass order for events. On the other hand, Koopman's approach to qualitative probability is not as restrictive and does yield a preference relation, which if suitably modified, is compatible with the natural subclass partial order.  In the work here, it is shown that a qualitative concept of conditional probability can be established on a thoroughly rigorous algebraic basis which can be directly related to Koopman qualitative conditional probability. In addition, it is shown that conditional events - which <u>must</u> be certain principal ideal cosets of events from the initial boolean algebra of events - can be compared, contrasted, and combined by any boolean operations, even if the events have distinctly different antecedents. In addition, conditional events form a conditional event algebra which in structure is midway between a pseudocomplemented distributive lattice and a full boolean algebra. This richness of structure allows for the development of an associated sound and complete conditional probability logic of propositions; deep structural relations with three-valued logic and DeFinetti's previous concept of partial indicator functions; and a novel interpretation of bayesian analysis from a qualitative/syntactic viewpoint.					
Published in <i>Proceedings of the 28th Annual Bayesian Research Conference</i> , 1989.					
14. SUBJECT TERMS  data fusion                      uncertainty measures combination of evidence      game theory					
17. SECURITY CLASSIFICATION OF REPORT  UNCLASSIFIED		18. SECURITY CLASSIFICATION OF THIS PAGE  UNCLASSIFIED		19. SECURITY CLASSIFICATION OF ABSTRACT  UNCLASSIFIED	
				16. PRICE CODE	
				20. LIMITATION OF ABSTRACT  SAME AS REPORT	

✓ UNCLASSIFIED

21a. NAME OF RESPONSIBLE INDIVIDUAL I. R. Goodman	21b. TELEPHONE (include Area Code) (619) 553-4014	21c. OFFICE SYMBOL Code 421

CONDITIONAL EVENT ALGEBRAS: TWO NEW CHARACTERIZATIONS AND THEIR  
RELATIONS TO BAYESIAN ANALYSIS

Invited presentation to the 27<sup>th</sup> Annual  
Bayesian Research Conference, Sponsored  
by the University of Southern California,  
Los Angeles, CA, under the direction of  
Prof. Ward Edwards, Feb. 16, 17, 1989.

Abstract

The standard approach in developing conditional probability is a numerically-oriented one, although one notable exception has been efforts in the area of qualitative (comparitive-preference ordering) probability theory, as developed by Suppes, Domotor, Fishburn, and others. However, the latter aspect of probability requires, in general, the preference ordering to be equivalent to the probability ordering. But, this is obviously too restrictive to be compatible with the basic monotonicity property of probability, relative to the usual partial subclass order for events. On the other hand, Koopman's approach to qualitative probability is not as restrictive and does yield a preference relation, which if suitably modified, is compatible with the natural subclass partial order.

In the work here, it is shown that a qualitative concept of conditional probability can be established on a thoroughly rigorous algebraic basis which can be directly related to Koopman qualitative conditional probability. In addition, it is shown that conditional events - which must be certain principal ideal cosets of events from the initial boolean algebra of events - can be compared, contrasted, and combined by any boolean operations, even if the events have distinctly different antecedents. In addition, conditional events form a conditional event algebra which in structure is midway between a pseudocomplemented distributive lattice and a full boolean algebra. This richness of structure allows for: the development of an associated sound and complete conditional probability logic of propositions; deep structural relations with three-valued logic and DeFinetti's previous concept of partial indicator functions; and a novel interpretation of bayesian analysis from a qualitative/syntactic viewpoint.



Accession for  
NTIS GRA&I ☒  
DTIC TAB ☐  
Unannounced ☐  
Justification

By  
Distribution/  
Availability Codes

Dist Avail and/or  
Special

A-1